

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 401 747  
A2**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 90110612.0

(51) Int. Cl.<sup>5</sup>: **A61K 31/445, A61K 31/52,  
A61K 31/57, A61K 37/02,  
A61K 31/71**

(22) Date of filing: 05.06.90

(30) Priority: 06.06.89 US 362354

(43) Date of publication of application:  
12.12.90 Bulletin 90/50

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **Calne, Roy**  
**22 Arrow Road**  
**Cambridge CB2 2AS(GB)**

(72) Inventor: **Calne, Roy**  
**22 Arrow Road**  
**Cambridge CB2 2AS(GB)**

(74) Representative: **Patentanwälte Grünecker,  
Kinkeldey, Stockmair & Partner**  
**Maximilianstrasse 58**  
**D-8000 München 22(DE)**

(54) **Methods of inhibiting transplant rejection in mammals using rapamycin and derivatives and prodrugs thereof.**

(57) This invention provides a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal a transplant rejection inhibiting amount of rapamycin. Also disclosed is a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal (a) an amount of rapamycin in combination with (b) an amount of one or more other chemotherapeutic agents for inhibiting transplant rejection, e.g., azathioprine, corticosteroids, cyclosporin and FK506, said amounts of (a) and (b) together being effective to inhibit transplant rejection and to maintain inhibition of transplant rejection.

**EP 0 401 747 A2**

# METHODS OF INHIBITING TRANSPLANT REJECTION IN MAMMALS USING RAPAMYCIN AND DERIVATIVES AND PRODRUGS THEREOF

## Field of the Invention

This invention relates to methods of inhibiting organ or tissue transplant rejection in mammals. More particularly, the invention relates to methods of inhibiting transplant rejection in mammals in need thereof, comprising administering to such mammals a transplant rejection inhibiting amount of rapamycin.

## Background of the Invention

Rejection and infective complications resulting from immunosuppressive treatment are the principal causes of failure of organ allografting in man, that is, an organ graft made between two genetically different individuals in the same *Homo sapiens* species. In order to minimize the individual specific side-effects of the three effective agents used in clinical practice, namely azathioprine, corticosteroids and cyclosporin, small doses of each are used in combination "triple therapy". Of the three agents currently used in such triple therapy, cyclosporin is the most powerful, but has the unsatisfactory side-effect of nephrotoxicity in man which can lead to structural renal damage. Increased corticosteroid dosage and antilymphocyte antibody preparations, poly- or monoclonal, are used for the treatment of rejection crises. A number of studies have been taken to investigate other potentially effective compounds for use as immunosuppressants and transplant rejection inhibitors, but to date, none have been found to be useful in clinical settings because of side-effects, such as toxicity, the lack of efficacy or a combination of these factors.

The fungal product FK506 was reported to have immunosuppressive activity in animals with organ grafts (Ochiai, T., et al., Transplant. Proc., Vol. XX, No. 1, pp. 209-214, 1988). Although the immunosuppressive activity of FK506 was confirmed, the toxicity in mammals, such as rats, pigs and dogs, and in primates, e.g., baboons, was too severe to proceed to clinical phase trials (Collier, D. St. J., et al., Transplant. Proc., Vol. XX, No. 1, pp. 226-228, 1988).

It would be extremely useful to discover a compound having immunosuppressive activity which could be employed to increase transplant acceptance in a recipient but without causing serious toxic side effects typically associated with conventional immunosuppressant therapy, such as those discussed above.

Rapamycin is a lipophilic macrolide with certain structural similarities to FK506 produced by *Streptomyces hygroscopicus* with both antifungal and antitumor properties (Sehgal, S.N. et al., J. Antibiot., Vol. 28, pp. 727-732, 1975; Eng. C.P., et al., J. Antibiot., Vol. 37, pp. 1231-1237, 1984).

It was reported that rapamycin inhibited two experimental immunopathies, i.e., experimental allergic encephalitis and adjuvant arthritis, and the formation of humoral (IgE-like) antibody. (Martel, R.R., et al., Can. J. Physio. Pharmacol., 55: 48-51, 1977). It has also been reported recently that rapamycin inhibits murine T cell activation, apparently through a different mechanism from both FK506 and cyclosporin. Thus, both FK506 and cyclosporin were found to act at an early stage of the immune response by blocking the formation of IL-2. Rapamycin, on the other hand, did not block the formation of IL-2 but acted at a later stage of the immune response by inhibiting the response of T cells to growth-promoting lymphokines. (Staruch, M.J., et al., The FASEB Journal, Vol 3, No. 3, abstract #3411, 1989). In addition, it was disclosed that rapamycin blocks the immunosuppressive effect of FK506 but not that of cyclosporin A (Dumont, F.J. et al., The FASEB Journal, Vol. 3, No. 4, abstract #5256, 1989). There was no teaching or suggestion in these reports, however, that rapamycin could or should be used to effectively inhibit organ or tissue transplant rejection in mammals. Furthermore, these reports do not disclose or intimate that the toxic side-effects associated with FK506, and other immunosuppressive agents, would not likewise arise from administering rapamycin as an agent to inhibit transplant rejection in transplant operations.

## Objects of the Invention

It is an object of this invention to provide a method of increasing allograft acceptance (or inhibiting organ or tissue transplant rejection) in mammals by administering an efficacious compound of low toxicity.

It is another object of this invention to reduce the toxicity of other conventional chemotherapeutic agents for inhibiting transplant rejection by combining their administration with an efficacious compound of low toxicity.

These and other objects of the invention will become clearer in light of the detailed description which follows.

## 5 Summary of the Invention

The present inventor has discovered a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal a transplant rejection inhibiting amount of rapamycin.

10 The present inventor has also discovered a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal (a) an amount of rapamycin in combination with (b) an amount of one or more other chemotherapeutic agents for inhibiting transplant rejection, said amounts of (a) and (b) being effective to inhibit transplant rejection and to maintain inhibition of transplant rejection.

15

## Detailed Description of the Invention

All patents and literature references are hereby incorporated by reference in their entirety.

20 The present invention provides a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal a transplant rejection inhibiting amount of rapamycin.

The present invention also provides a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal an amount of rapamycin effective to 25 inhibit transplant rejection and to maintain inhibition of transplant rejection.

As used herein, the terms "inhibiting organ or tissue transplant rejection" and "maintain inhibition of transplant rejection" refer to increasing organ or tissue transplant acceptance (or decreasing the likelihood of organ or tissue transplant rejection) involving allografts, i.e., transplantation of organs or tissues from donor to recipient both of whom are in the same species (intraspecific), such as Homo sapiens.

30 Rapamycin is an antifungal antibiotic which is extractable from a streptomycete, e.g., Streptomyces hygroscopicus. Methods for the preparation of rapamycin are disclosed in Sehgal et al., U.S. Patent Nos. 3,929,992, and 3,993,749. In addition, monoacyl and diacyl derivatives of rapamycin and methods for their preparation are disclosed by Rakhit, U.S. Patent No. 4,316,885. Furthermore, Stella et al., U.S. Patent No. 4,650,803 disclose water soluble prodrugs of rapamycin, i.e., rapamycin derivatives including the following 35 rapamycin prodrugs: glycinate prodrugs, propionate prodrugs and the pyrrolidino butyrate prodrugs.

The methods and compositions of the present invention include the use of natural and synthetic rapamycin, genetically engineered rapamycin and all derivatives and prodrugs of rapamycin, such as described in the aforementioned U.S. patents, U.S. Patent Nos. 3,929,992; 3,993,749; 4,316,885; and 4,650,803, the contents of which are hereby incorporated by reference.

40 The present inventor has noted the efficacy of rapamycin in inhibiting transplant rejection, e.g., by depressing the immune system in mammals without the attendant toxic side-effects associated with other conventional immunosuppressive agents, e.g., azathioprine, corticosteroids and cyclosporin. Among such toxic side-effects are nephrotoxicity, severe leukopenia, thrombocytopenia, Cushing's Syndrome and diabetes.

45 It has been discovered that rapamycin reduces or inhibits allograft rejection in mammals, i.e., organ or tissue transplantation from donor to recipient of the same species. Among such transplanted organs or tissues and given illustratively, are heart, liver, kidney, spleen, lung, small bowel, pancreas, skin, and bone marrow, or a combination of any of the foregoing.

50 As used herein, the term "transplant rejection inhibiting amount" refers to the amount of rapamycin (or of rapamycin in combination with one or more other chemotherapeutic agents for inhibiting transplant rejection) which may be administered so as to inhibit transplant rejection in a mammal and to maintain transplant rejection inhibition, without causing severe toxic side-effects, e.g., nephrotoxicity, renal failure, etc. Those skilled in the art will appreciate that the dosage or amount of a transplant rejection inhibiting compound which is administered to a subject about to undergo or having undergone an organ or tissue 55 transplant, will vary according to a number of factors, including individual characteristics, such as weight, age, and other factors, such as the type of organ or tissue transplanted or about to be transplanted.

In one aspect of this invention, a tissue rejection inhibiting amount of rapamycin comprises from about 0.5 to about 50 mg/kg/day, preferably from about 1 to about 5 mg/kg/day. Further studies indicate that an

effective therapeutic dose of rapamycin for inhibition of rejection comprises about 0.01 to about 10 mg/kg/day, and preferably about 0.025 to about 5 mg/kg/day. Such doses may be given intermittently, for example, every other day or every third day. When administered in combination with one or more other chemotherapeutic agents for inhibiting rejection in organ or tissue transplant, the effective therapeutic dose of rapamycin may be even less than indicated above. In another aspect, the inhibiting transplant rejection amount of rapamycin is administered for a period of time comprising from about 1 to about 180 days, or longer, as necessary. Those skilled in the art will recognize that compounds, drugs, agents, and the like, for inhibiting transplant rejection, may be administered to a subject mammal, e.g., a human, for an indefinite post-transplantation period, in some instances, for the lifetime of the subject, provided, of course, that the subject is tolerating the compound, drug, agent, etc., reasonably well without serious side-effects.

Rapamycin may be administered either orally or parenterally, e.g., intramuscularly, intraperitoneally, subcutaneously, or intravenously to a mammal subject. The preferred route of administration is oral.

According to this invention, rapamycin may be administered in various pharmaceutical forms, including pharmaceutical forms suitable for parenteral injectable use, such as sterile aqueous solutions or dispersions and sterile powders for the preparation of sterile injectable solutions or dispersions. In addition, rapamycin may be administered in tablets, caplets, capsules, and the like for convenient oral administration. Rapamycin may be administered in a pharmaceutically compatible or acceptable carrier, which includes by way of non-limiting example, oils, e.g., olive oil, alcohols, propylene glycol and polyethylene glycols, and surfactants, such as Chemophor EL (BASF), and polysorbate 80.

Another useful feature of this invention resides in the administration of rapamycin in combination with other conventional drug therapies, such as "triple therapy," azathioprine (available from Burroughs Wellcome Co., Research Triangle Park, N.C., under the tradename Imuran®), corticosteroids (available from the Upjohn Company, Kalamazoo, Michigan, under the tradename Solu-Medrol®); cyclosporin (and cyclosporin A) (available from Sandoz Pharmaceuticals, East Hanover, New Jersey, under the tradename Sandimmune®), and also FK506, (available from Fujisawa Pharmaceutical Co., Ltd., Osaka, Japan, under the tradename Fujimycin®). By combining rapamycin with such other conventional chemotherapeutic drugs or agents for inhibiting transplant rejection, the toxicity of the latter may be advantageously reduced in that lesser amounts of such toxic drug or agents are required in order to inhibit transplant rejection in a mammalian transplant subject either before or after transplantation, and also to maintain inhibition of transplant rejection.

Thus, the present invention provides a method of inhibiting transplant rejection in a mammal in need thereof, comprising administering to said mammal (a) an amount of rapamycin in combination with (b) an amount of one or more other chemotherapeutic agents for inhibiting transplant rejection, the amounts of (a) and (b) together being effective to inhibit transplant rejection and to maintain inhibition of transplant rejection. It should be understood that the amount of either (a) or (b) alone, might or might not be effective in inhibiting and maintaining transplant rejection. The combination of the two components (a) and (b), together, however, is effective to inhibit transplant rejection and to maintain inhibition of transplant rejection.

Therefore, as used herein, such "other" chemotherapeutic agents for inhibiting transplant rejection include, for example, azathioprine, corticosteroids, cyclosporin (and cyclosporin A) poly and monoclonal antilymphocyte antibodies (OKT3) and FK506, or a combination of any of the foregoing.

The various features of the invention described above, such as the type of organ or tissue which is transplanted; the amount effective to inhibit transplant rejection and to maintain inhibition of transplant rejection; the mode or route of administration; and the duration of treatment; apply to the method of inhibiting organ or tissue transplant rejection by administering rapamycin in combination with one or more other chemotherapeutic agents for inhibiting transplant rejection and to maintain inhibition of transplant rejection.

In addition, however, it should be understood that such other chemotherapeutic agents may be administered continuously or intermittently with rapamycin. Furthermore, the route of administration may differ from that employed for rapamycin. That is to say, such other chemotherapeutic agents may be administered parenterally while rapamycin is being administered orally to the mammalian subject.

The present invention is described below in specific working examples which are intended to illustrate the invention without limiting its scope.

## 55 EXAMPLE 1: ORGAN ALLOGRAFT STUDIES

### Rats:

Heterotopic heart allografts were performed from DA donor to PVG recipients. DA and PVG refer to specific rat strains. Rapamycin was administered intramuscularly in olive oil for the first 10 postoperative days except for Group 6 which only received the drug on days 3-6 postoperatively. Graft survival was assessed by daily palpation.

#### Large Animals:

On the basis of the initial toxicity study two experiments were considered suitable to test whether rapamycin was immunosuppressive, namely, a short term dosing in dogs following renal transplantation and indefinite dosing of pigs, again following renal grafting.

#### Pigs:

In pigs an orthotopic renal transplantation model was used and Mixed Lymphocyte Cultures (MLC) were performed to confirm histoincompatibility between donor and recipient. Untreated controls mean survival time is less than 10 days. Rapamycin was administered at a dose of 2mg/kg orally daily commencing on the first postoperative day.

#### Results: Allograft Studies

#### Rat:

For the rat, survival of allografts is shown in Table 1.

TABLE 1

SURVIVAL OF HETEROTOPIC CARDIAC ALLOGRAFTS IN RATS			
Group 1	No. Rats	Dose Schedule	Survival (days)
1	4	50mg/kg x 10d	75*, 88*, 100(x2)
2	4	10mg/kg x 10d	65, 77, 88, 100
3	4	2mg/kg x 10d	58, 59, 59, 66
4	4	1mg/kg x 10d	34, 49, 52, 55
5	4	0.5mg/kg x 10d	19, 20, 20, 35
6	5	10mg/kg d3-6	15, 19, 19, 19, 21
N.B. Rapamycin was administered intramuscularly in olive oil. (18mg/ml and 10mg/ml suspensions used.)			
Rat strains used: DA donors in PVG recipients.			
control rejection time (n = 10) = 7.4 days.			

Explanation of Table 1: Heterotopic heart allografts in the neck of the rats were performed from DA donors to PVG recipients using the surgical techniques previously described (Heron, I., *Acta, Pathol. Microbiol. Scand.* 79:366, 1971). Rapamycin was dissolved in olive oil at a maximum concentration of 15 mg/ml and administered by daily intramuscular injection at dose schedules varying from 0.5 mg/kg to 50 mg/kg for ten consecutive days and in the last group at 10 mg/kg on days 3 to 6. Graft survival was assessed by daily palpation of the heart.

Rapamycin prolonged allograft survival at all doses tested. Although there was some loss of weight this was not as marked as that found when FK506 was administered to rats.

Dogs

In dogs dosing at all levels induced a vasculitis and at doses greater than 0.25 mg/kg this led to such severe manifestations that they were killed before the end of the 28 day study. At the higher doses the vasculitis affected the gastrointestinal tract, interestingly, it also caused a thrombocytopaenia. Marked depletion of cells in the lymphoid tissue, particularly B cells, occurred. In the dog, toxicity due to a vasculitis, that appears to have a particular predilection for the gastrointestinal tract, made it impossible to assess the immunosuppressive effects of the drug in this particular model. This species-specific reaction to rapamycin confirmed similar unpublished observations of the inventor and his associates.

Pigs

Survival and cause of death are shown in Table 2 with current creatinine values.

TABLE 2

SURVIVAL OF PIG ALLOGRAFTS RECEIVING RAPAMYCIN 2mg/Kg/DAY ORALLY IN DAYS AND CURRENT CREATININE						
	SURVIVAL	RENAL HISTOLOGY	CREATININE	OUTCOME	CAUSE OF DEATH	DAY LAST DOSED
1	> 51		217	ALIVE		43
2	> 66		193	ALIVE		52
3	> 72		283	ALIVE		64
4	4	ACUTE REJECTION	1,410	DIED	REJECTION	4
5	5	INFARCTION		DIED	INFARCTION	5
6	48	FOCI PYELONEPHRITIS	161	DIED	INTERSTITIAL PNEUMONITIS	47
7	49	MILD ATN	235	DIED	INTERSTITIAL PNEUMONITIS	48
8	50	MILD ATN	176	DIED	INTERSTITIAL PNEUMONITIS	49
9	55	EARLY MILD REJECTION	239	DIED	INTERSTITIAL PNEUMONITIS	51
10	63	MODERATE ATN	283	DIED	INTERSTITIAL PNEUMONITIS	62

Explanation of Table 2: Orthotopic kidney transplantation with contralateral nephrectomy was performed in the pig, as previously described (Calne, R.Y. et al., Brit. J. Surg., 59: 969-977 (1972). Donor and recipient pairs were obtained from litters with different parents and incompatibility at the Major Histocompatibility Complex (MHC) was confirmed by the mixed lymphocyte reaction (Bradley, B.A., et al., Tissue Antigens, 4: 283-290, 1974). Rapamycin was administered orally at 2 mg/kg/day dissolved in olive oil at a concentration of 10 mg/ml.

In the case of pigs, one died of accelerated acute rejection and one died due to technical failure. The remaining eight animals recovered well and after an initial weight loss of approximately 10%. Subsequently, at about day 50 5 animals developed anorexia, diarrhea and became unwell to the extent that it was decided that they should be killed. Histological examination of these animals revealed that they were suffering from interstitial pneumonitis, probably due to over-immunosuppression, and this was the reason that they became unwell. Furthermore, the renal histology did not show evidence of rejection except a mild degree in one animal who had not received the drug for 4 days. Histological examination of the colons in these animals showed mucosal and submucosal edema but no vasculitis and no ulceration. Thus this was probably secondary to the systemic effects of the pneumonitis. The remaining three animals continued to thrive, all dosing having being stopped as indicated in Table 3.

## Discussion

Rapamycin was immunosuppressive and not toxic in the rat down to a dose of 0.5mg/kg although the compound was more effective at higher doses.

In pigs, the results of the toxicity study showed that the drug was tolerated at a dose of 1 mg/kg in that both animals gained weight. On histological examination colitis was seen but no vasculitis was found or suggested. Rapamycin was effective as an immunosuppressive agent but after about 50 days of continuous dosing at 2 mg/kg, 50% of the animals developed interstitial pneumonitis due to over-immunosuppression, and these animals were killed. However, none of the animals showed any evidence of ulceration in the colon or vasculitis. Therefore, in future studies monitoring of blood drug levels will be of benefit.

In conclusion, rapamycin is a very effective immunosuppressive agent which can be employed to inhibit allograft transplantation rejection in mammalian subjects.

## Claims

1. A composition for inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising rapamycin for administration to said mammal in a transplant rejection inhibiting amount.

2. A composition according to claim 1, wherein the amount of rapamycin further is effective to maintain inhibition of transplant rejection.

3. A composition according to claim 1 or 2, comprising a combination of (a) an amount of rapamycin (b) an amount of one or more chemotherapeutic agents for inhibiting transplant rejection, said amounts of (a) and (b) together being effective to inhibit transplant rejection and to maintain inhibition of transplant rejection.

4. The composition according to claim 3, wherein said chemotherapeutic agent is selected from the group consisting of azathioprine, corticosteroids, cyclosporin and FK506 and combinations of any of the foregoing.

5. The composition according to any one of claims 1 to 4, wherein said transplant rejection inhibiting amount comprises from about 0.5 to about 50 mg/kg/day.

6. The composition according to any one of claims 1 to 4, wherein said amount comprises from about 1 to about 5 mg/kg/day.

7. The composition according to claim 5, wherein said amount comprises from about 0.01 to about 10 mg/kg/day.

8. The composition according to claim 7, wherein said amount comprises from about 0.025 to about 5 mg/kg/day.

9. The composition according to any one of claims 1 to 8, wherein said transplant rejection inhibiting amount of rapamycin is administered to said mammal for a period of time comprising from about 1 to about 180 days.

10. The composition according to any one of claims 1 to 9, wherein said rapamycin is formulated in an



oral dosage form.

11. The composition according to any one of claims 1 to 9, wherein said rapamycin is formulated in a parenteral dosage form.

12. Use of rapamycin for the preparation of a medicament for inhibiting organ or tissue transplant rejection in a mammal in need thereof.

13. Use of rapamycin according to claim 12, wherein the medicament further serves to maintain said inhibition of organ or tissue transplant rejection.

14. Use of rapamycin according to claim 12 or 13, wherein the medicament further contains an amount of one or more other chemotherapeutic agents for inhibiting organ or tissue transplant rejection in a mammal in need thereof.

15. Use of rapamycin according to claim 14, wherein said chemotherapeutic agent is selected from the group consisting of azathioprine, corticosteroids, cyclosporin, FK506, OKT3 and combinations of any of the foregoing.

16. Use of rapamycin according to any one of claims 12 to 15, wherein said medicament is formulated for administration of rapamycin to said mammal in an amount of from about 0.5 to about 50 mg/kg/day.

17. Use of rapamycin according to any one of claims 12 to 15, wherein said amount comprises from about 1 to about 5 mg/kg/day.

18. Use of rapamycin according to claim 16, wherein said amount comprises from about 0.01 to about 10 mg/kg/day.

19. Use of rapamycin according to claim 18, wherein said amount comprises from about 0.025 to about 5 mg/kg/day.

20. Use of rapamycin according to any one of claims 12 to 19, said medicament being administered to said mammal for a period of time comprising from about 1 to about 180 days.

21. The use of rapamycin according to any one of claims 12 to 20, wherein said rapamycin is administered in an oral dosage form.

22. The use of rapamycin according to any one of claims 12 to 20, wherein said rapamycin is administered in a parenteral dosage form.

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 401 747 A3**

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 90110612.0

(51) Int. Cl.<sup>5</sup>: **A61K 31/445, A61K 31/52,**  
**A61K 31/57, A61K 37/02,**  
**A61K 31/71**

(22) Date of filing: 05.06.90

(30) Priority: 06.06.89 US 362354

(43) Date of publication of application:  
12.12.90 Bulletin 90/50

(64) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU NL SE**

(69) Date of deferred publication of the search report:  
03.07.91 Bulletin 91/27

(71) Applicant: **Calne, Roy**  
**22 Arrow Road**  
**Cambridge CB2 2AS(GB)**

(72) Inventor: **Calne, Roy**  
**22 Arrow Road**  
**Cambridge CB2 2AS(GB)**

(74) Representative: **Patentanwälte Grünecker,**  
**Kinkeldey, Stockmair & Partner**  
**Maximilianstrasse 58**  
**W-8000 München 22(DE)**

(54) **Methods of inhibiting transplant rejection in mammals using rapamycin and derivatives and prodrugs thereof.**

(57) This invention provides a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal a transplant rejection inhibiting amount of rapamycin. Also disclosed is a method of inhibiting organ or tissue transplant rejection in a mammal in need thereof, comprising administering to said mammal (a) an amount of rapamycin in combination with (b) an amount of one or more other chemotherapeutic agents for inhibiting transplant rejection, e.g., azathioprine, corticosteroids, cyclosporin and FK506, said amounts of (a) and (b) together being effective to inhibit transplant rejection and to maintain inhibition of transplant rejection.

**EP 0 401 747 A3**



European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 90 11 0612

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X,D	CAN. J. PHYSIOL. PHARMACOL., vol. 55, no. 1, February 1977, pages 48-51; R.R. MARTEL et al.: "Inhibition of the immune response by rapamycin, a new antifungal antibiotic" * Whole article *	1,2,5-10, 12,13, 16-21	A 61 K 31/445 A 61 K 31/52 A 61 K 31/57 A 61 K 37/02 A 61 K 31/71
X,D	FASEB J., vol. 3, no. 4, 1989, 73rd Ann. Meet. Fed. Am. Soc. Exp. Biol., New Orleans, 19th - 23rd March 1989, page A1131, abstract no. 5256; F.J. DUMONT et al.: "Rapamycin blocks the immunosuppressive effect of FK506 but not that of cyclosporin A" * Whole abstract *	1,3,4,12, 14,15	
Y,D	INTERNATIONAL ORGAN TRANSPLANT FORUM, Pittsburg, Pennsylvania, 8th - 11th September 1987, vol. 20, no. 1, suppl. 1, pages 209-214; T. OCHIÁI et al.: "Studies on FK506 in experimental organ transplantation" * Whole article *	3-8,14, 15,17	
P,X	PROGRESS IN IMMUNOLOGY, 7TH INTERNATIONAL CONGRESS OF IMMUNOLOGY, Berlin, 1989, vol. VII, pages 1195-1198; B.M. MEISER et al.: "Rapamycin: A new and highly active immunosuppressive macrolide with an efficacy superior to cyclosporine" * Whole article *	1-22	
P,X	4TH CONGRESS OF THE EUR. SOC. ORGAN TRANSPLANTATION, Barcelona, 1st - 4th November 1989, TRANSPLANT. PROC., vol. 22, no. 4, August 1990, pages 1638-1641, R.E. MORRIS et al.: "A study of the contrasting effects of cyclosporine, FK 506, and rapamycin on the suppression of allograft rejection" * Whole article *	1,2,9-13, 16,18-22	
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		13 March 91	GAC G.
<b>CATEGORY OF CITED DOCUMENTS</b> X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			



European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 90 11 0612

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
Y	IDEM	3-8,14, 15,17	
P,X	THE LANCET, vol. 2, no. 8656, 22nd July 1989, page 227; R.Y. CALNE et al.: "Rapamycin for immunosuppression in organ allografting" * Whole article *	1,2,9, 11-13,16, 18-20,22	
Y	IDEM	3-8,14, 15,17	
P,X	MED. SCI. RES., vol. 17, no. 20, 16th - 31st October 1989, pages 609-610; R.E. MORRIS et al.: "Identification of a new pharmacologic action for an old compound" * Whole article *	1,2,9,10, 12,13,16, 18-21	
Y		3-8,14, 15,17	
A,D	THE JOURNAL OF ANTIBIOTICS, vol. 37, no. 10, 1984, pages 1231-1237; C.P. ENG et al.: "Activity of rapamycin (AY-22,989) against transplanted tumors" * Whole article *	1,2,5-13, 16-22	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		13 March 91	GAC G.
<b>CATEGORY OF CITED DOCUMENTS</b> X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention		E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document	